

# What's an Ionic Liquid?

by **Keith E. Johnson**

*"When I use a word," Humpty Dumpty said, in rather a scornful tone,  
"it means just what I choose it to mean – neither more nor less."*

*"The question is," said Alice, "whether you can make words mean so many different things."*

LEWIS CARROLL, *Through the Looking Glass*

## Introduction to the Phrase "Ionic Liquids"

**T**he Structure and Properties of Ionic Melts" was the title of a Faraday Society Discussion held in Liverpool in 1961; it dealt exclusively with molten inorganic salts.<sup>1</sup> "Ionic Liquids" was the title of Chapter 6 of the textbook *Modern Electrochemistry* by Bockris and Reddy, published in 1970: it discussed liquids ranging from alkali silicates and halides to tetraalkylammonium salts.<sup>2</sup> The modern era of ionic liquids stems from the work on alkyipyridinium and dialkylimidazolium salts in

scheduling of groups of papers. EUCHEM 2006 dealt with Molten Salts and Ionic Liquids with the content of each session usually mixed.

## Modern Ionic Liquids

The properties of a modern ionic liquid are summarized in Table I. Particularly significant are (i.) the low vapor pressures in most instances which contrast the environmental problems of volatile organic solvents and (ii.) moderate specific conductivities, usually in the same range as

One property that we emphasized recently is the molarity of the liquid<sup>6</sup> (a straightforward quantity except for mixed systems such as a basic chloroaluminate containing both Cl<sup>-</sup> and AlCl<sub>4</sub><sup>-</sup> in significant amounts). The molarity is important regarding kinetic measurements, including conductivities.

Table II indicates a range of molarities of many liquids from 1 to 60, with water at 55, liquid alkali halides up to 35 (LiCl) and most organic salts less than 10. Specific conductivities span a far greater range from the metal sodium through molten inorganic salts in the Scm<sup>-1</sup> region to organic salts (the modern ionic liquids) and aqueous solutions in the mScm<sup>-1</sup> region and finally to the near non-conducting but ionizing acetic acid and water at μScm<sup>-1</sup>. Combining these data into molar conductances is illuminating. We see comparable values for simple inorganic salts alone and in aqueous solutions\* but much smaller values for the low temperature semi-organic and organic systems. Thus these modern ionic liquids must consist of IONS AND ION PAIRS, (undissociated molecules), while liquid alkali halides are purely IONIC and aqueous electrolytes behave as a mixture of hydrated ions and the molecular solvent water. Figure 1 attempts to picture these differences.

## Acid-Base Properties and Water Interactions

While simple salts such as KCl can be thought of as the product of an electron transfer between elements, organic salts can be traced to a proton transfer between an acid and base. Cations such as emim<sup>+</sup> and n-bupy<sup>+</sup> result from the alkylation of the bases

\* Extrapolation of aqueous solution molar or equivalent conductances to infinite dilution, at which ion pairing is eliminated, gives many values in the 100 to 150 Scm<sup>2</sup> mol<sup>-1</sup> range,<sup>11</sup> the exceptions involving H<sup>+</sup> and OH<sup>-</sup> for which the Grotthus mechanism operates.<sup>2</sup>

Table I. Modern ionic liquids.

|                         |  |
|-------------------------|--|
| A salt                  | Cation and or anion quite large                      |
| Freezing point          | Preferably below 100°C                               |
| Liquidus range          | Often > 200°C  |
| Thermal stability       | Usually high   |
| Viscosity               | Normally < 100 cP, workable                          |
| Dielectric constant     | Implied ≤ 30   |
| Polarity                | Moderate   |
| Specific conductivity   | Usually < 10 mScm <sup>-1</sup> , "Good"             |
| Molar conductivity      | < 10 Scm <sup>2</sup> mol <sup>-1</sup>              |
| Electrochemical window  | > 2V, even 4.5 V, except for Brønsted acidic systems |
| Solvent and/or catalyst | Excellent for many organic reactions                 |
| Vapor pressure          | Usually negligible                                   |

Colorado in the late 1970s.<sup>3</sup> The term ionic liquids was introduced<sup>4</sup> to cover systems below 100°C, one reason being to avoid the words "molten salts" in phrases such as "ambient temperature molten salts," another to create an impression of freshness and a third, perhaps, for patent purposes. The first "Conference on Ionic Liquids" took place in Salzburg in 2005, Molten Salts 7 in Toulouse in 2005 had one of ten sessions devoted to Ionic Liquids; but the International Symposia on Molten Salts of ECS from 1976 to the present have not shown discrimination on the basis of temperature, beyond

those of aqueous electrolytes. It is found that many such systems are excellent solvents or catalysts for organic reactions<sup>3</sup> and some simple processes such as electrodeposition.<sup>5</sup> Unfortunately, one finds reports of new "ionic liquids" without data on conductivities, which would establish that they are dissociated to some extent at least into ions.


## Liquids Comparisons

How do these ionic liquids compare with other liquids, especially those which conduct electricity? Table II presents some illustrative data.

Table II. Properties of various liquids.

| Compound  | Temp. °C | Molarity Mol L <sup>-1</sup> | Spec Cond Scm <sup>-1</sup> | Molar Cond Scm <sup>2</sup> mol <sup>-1</sup> | Visc. cP | Ref. |
|---|----------|------------------------------|-----------------------------|---|----------|------|
| <i>Ionic Liquids</i>  |          |                              |                             |   |          |      |
| NaCl  | 900      | 25.3                         | 3.88                        | 154   | 1.05     | 2    |
| LiCl  | 780      | 35                           | 7.59                        | 217   |          | 2    |
| NaNO <sub>3</sub>   | 450      | 21.4                         | 0.72                        | 34  | 6.9      | 2    |
| Na <sub>2</sub> SiO <sub>3</sub>  | 1750     | ≈ 18                         | 4.8                         | ≈ 270   |          | 2    |
| AlCl <sub>3</sub> (63%) – NaCl  | 175      | 5.8                          | 0.24                        | 41  | 3.85     | 7    |
| LiCl-KCl  | 450      | 29.7                         | 1.57                        | 53  | 2.44     | 7    |
| [(CH <sub>3</sub> ) <sub>3</sub> S][HBr <sub>2</sub> ]  | 25       | 7.3                          | 0.034                       | 4.7   | 20.5     | 8    |
| emimCl(60%)-AlCl <sub>3</sub>   | 25       | 5.3                          | 0.0065                      | 1.2   | 47       | 8    |
| emim Al <sub>2</sub> Cl <sub>7</sub>  | 25       | 3.4                          | 0.015                       | 4.4   | 14       | 8    |
| bmim CF <sub>3</sub> CO <sub>2</sub>  | 20       | 5.1                          | 0.0032                      | 0.6   | 73       | 8    |
| bupy BF <sub>4</sub>  | 25       | 5.5                          | 0.0019                      | 0.3   | 103      | 8    |
| emim[(CF <sub>3</sub> SO <sub>2</sub> ) <sub>2</sub> N]   | 25       | 3.7                          | 0.0057                      | 1.5   | 31       | 8    |
| <i>Others</i>   |          |                              |                             |   |          |      |
| H <sub>2</sub> O  | 25       | 55.3 <sup>a</sup>            | 4 x 10 <sup>-8</sup>        | 7 x 10 <sup>-7</sup>                          | 0.895    | 2    |
| 0.1 M aq. KCl   | 25       | 0.1 <sup>b</sup>             | 0.013                       | 129   | 0.9      | 2    |
| Na  | 100      | 40.4                         | 1.04 x 10 <sup>5</sup>      | 2.6 x 10 <sup>6</sup>                         | 1.058    | 2    |
| H <sub>2</sub> SO <sub>4</sub>  | 25       | 0.049 <sup>b</sup>           | 0.0104                      | 212   | 24.55    | 9    |
| CH <sub>3</sub> COOH  | 25       | 17.5 <sup>a</sup>            | 8 x 10 <sup>-9</sup>        | 4.6 x 10 <sup>-7</sup>                        | 1.056    | 10   |
| HF  | 0        | 50.1 <sup>a</sup>            | 1 x 10 <sup>-6</sup>        | 2 x 10 <sup>-5</sup>                          | 0.256    | 9    |
| a = total molarity • b = ionic molarity • emim = 1-ethyl-3-methyl-1 <i>H</i> -imidazolium •<br>bmim = 1-butyl-3-methyl-1 <i>H</i> -imidazolium • bupy = 1-butylpyridinium |          |                              |                             |   |          |      |

Hmim<sup>+</sup> and py<sup>+</sup>. Anions such as NO<sub>3</sub><sup>-</sup>, AlCl<sub>4</sub><sup>-</sup> and PF<sub>6</sub><sup>-</sup> derive from Lewis acid – base reactions:



It is also possible for Lewis neutral ions to add further Lewis acids to form acidic ions, *e.g.*, Al<sub>2</sub>Cl<sub>7</sub><sup>-</sup>, Sb<sub>2</sub>F<sub>11</sub><sup>-</sup>, HBr<sub>2</sub><sup>-</sup>.

As a result we can obtain Lewis basic, neutral or acidic ionic liquids, exemplified by the well-researched haloaluminates. In addition we can make protic acidic liquids which are simultaneously of any Lewis aspect.<sup>12</sup> How is proton chemistry in ionic

liquids related to proton chemistry in water?

First, acidic protons in ionic liquids often occur as anions, *e.g.*, HCl<sub>2</sub><sup>-</sup>, HBr<sub>2</sub><sup>-</sup>, H<sub>2</sub>Br<sub>3</sub><sup>-</sup>, rather than cations. Second, protons bonded to bases such as pyridine and 1-methylimidazole are not labile.<sup>13</sup> Do these bases behave in line with their p*K<sub>a</sub>*'s in water? No. Bases in ionic liquids appear to act in accordance with their gas phase proton affinities (1-methylimidazole > pyridine > ammonia).<sup>14</sup> The complicating factor of course is solvation in water. One can, however, obtain Hammett acidity data for some systems which make fair chemical sense.<sup>15</sup>

What happens when water is added to an ionic liquid or not completely removed at some stage in its preparation?

1. Some anions such as AlCl<sub>4</sub><sup>-</sup> and HCl<sub>2</sub><sup>-</sup> are irreversibly decomposed.
2. Water may bind strongly to one of the ions.
3. Water may dissolve the liquid until it forms a saturated salt solution; unlike the case of dissolving a solid salt, the other phase will be liquid and it may or may not contain water.
4. At high temperatures, species such as Li(H<sub>2</sub>O)<sub>4</sub><sup>+</sup>Cl<sup>-</sup> may decompose to LiOH and HCl<sup>7</sup>.

(continued on next page)




Fig. 1. Representation of (a) liquid NaCl, (b) molten bmim PF<sub>6</sub>, and (c) aqueous NaCl. Courtesy of A. East and Q. Li

Table III. Electrode processes of ionic liquid solvents.

| Process             | Examples   |
|---------------------|--|
| 1. Cation Reduction | $K^+ + e^- \rightarrow K$<br>$bmim^+ + e^- \rightarrow bmim \rightarrow bmim_2 \rightarrow$  |
| 2. Anion Reduction  | $NO_3^- + 2e^- \rightarrow NO_2^- + O^{2-}$<br>$O^{2-} + NO_3^- \rightarrow NO_2^- + O_2^{2-}$<br>$CO_3^{2-} + 4e^- \rightarrow C + 3O^{2-}$<br>$Al_2Cl_7^- + 3e^- \rightarrow Al + AlCl_4^- + 3Cl^-$        |
| 3. Anion Oxidation  | $SO_4^{2-} - 2e^- \rightarrow SO_3 + \frac{1}{2} O_2$<br>$CO_3^{2-} - 2e^- \rightarrow CO_2 + \frac{1}{2} O_2$<br>$BF_4^- - e^- \rightarrow BF_3 + \frac{1}{2} F_2$<br>$F_2 + bmim^+ \rightarrow C_nF_{n+2}$ |
| 4. Proton Reactions | $NO_3^- + H_2O + 2e^- \rightarrow NO_2^- + 2OH^-$<br>$HCl_2^- + e^- \rightarrow \frac{1}{2} H_2 + 2Cl^-$   |

### Electrochemistry of Ionic Liquids— The General Situation

Once we have established from conductivity measurements that a liquid is ionic (not ionic/molecular), its temperature and complexity should not pose special problems for using it as a solvent for electrochemistry. Table III lists the types of electrode processes one encounters.<sup>16</sup> With inorganic cations, the solubility of a metal in its salt may arise while the reduction of organic cations is complex and irreversible. Following chemical reactions such as attack of an electrode reaction product on the opposite ion in the system can also occur ( $F_2$  vs.  $bmim^{+17}$ ,  $O^{2-}$  vs.  $NO_3^-$ ).

### Detecting Your Ionic Liquid

The following (Scheme 1) is a sequence of steps prompted by finding two 19<sup>th</sup> century pieces of work: (a) Gardner's observation<sup>18</sup> of maximum-boiling point compositions for mixtures of several bases with acetic acid (he incorrectly called these specific mixtures compounds not azeotropes); and (b) Zawidzki's vapor pressure studies<sup>19</sup> of pyridine mixed with formic, acetic and propionic acids (good data even if some fine details are questionable).

#### Scheme 1. Detecting an ionic liquid.

(a) For an acid-base system  $HA + B$ :

**Clue #1** Vapor pressures of some mixtures show large negative deviation from Raoult's law: compound forms

**Clue #2** <sup>1</sup>H NMR spectra show downfield shift of HA's proton with B addition: BHA is ionized to  $BH^+A^-$

**Clue #3** Some mixtures show  $\sigma$  values in the  $mScm^{-1}$  range, far greater than for the components:  $BH^+A^-$  dissociates

The overall chemistry



is then governed by the ionization and dissociation constants

$$K_i = \frac{[BH^+][A^-]}{[B][AH]}$$

and

$$K_d = \frac{[BH^+][A^-]}{[BH^+A^-]}$$

with the ionic liquid containing the ions  $BH^+$  and  $A^-$  and the ion-pair  $BH^+A^-$ .<sup>20-22</sup> In practice, one may also have to include the self-ionization of HA and the formation of hydrogenates  $H_nA_{n+1}$ .

(b) For the salt MX or BRA where R = alkyl:

**Clue #1** Very low vapor pressures: stable compound in liquid

**Clue #2** NMR data less dramatic: may suggest  $BR^+A^-$

**Clue #3**  $\sigma$  values at least in the  $mScm^{-1}$  range: MX or  $BR^+A^-$  dissociation

For the acid-base systems the vapor may consist of the ionic compound or the original components,<sup>23</sup> whereas the breakdown of  $BR^+A^-$  (e.g., emimCl) into B and RA is probably irreversible.<sup>24,25</sup> The behavior of the vapor should correlate with the gas phase proton affinities.

### Conclusion

An ionic liquid is characterized by a specific conductivity in the  $mScm^{-1}$  range as a minimum, together with a molar conductivity probably exceeding  $0.1 Scm^2 mol^{-1}$ . In addition, the liquid should only contain ions with lesser numbers of ion pairs or parent molecules. Ionic liquids tend to have low dielectric constants, which means they are not ionizing solvents. One can assign an ionicity to an ionic liquid in terms of its conductivity and, presumably one might attempt to describe any liquid in this manner but ignoring the chemical properties of excess solvent molecules could lead to significant errors.

Thus in answer to the opening question, "What is an ionic liquid?" I prefer the answer, "Just a liquid salt consisting of ions and ion pairs." ■

### Acknowledgments

The long-term support of NSERC (Canada) and recent contributions of the Petroleum Technology Research Centre (Regina) and the Saskatchewan Research Council are appreciated.

### References

1. The Structure and Properties of Ionic Melts, *Discuss. Faraday Soc.*, **32**, University Press, Aberdeen (1962).
2. J. O'M Bockris and A. K. N. Reddy, *Modern Electrochemistry*, Plenum, New York (1970).
3. J. S. Wilkes, in *ACS Symposium 818*, R. D. Rogers and K. R. Seddon, Editors, Washington, DC, p. 214 (2002); J. S. Wilkes, *J. Mol. Cat. A: Chemical*, **214**, 11 (2004).
4. K. R. Seddon, *J. Chem. Tech. Biotechnol.*, **68**, 351 (1997).
5. Q. Zhu and C. L. Hussey, *J. Electrochem. Soc.*, **149**, C268 (2002).
6. L. Xiao, J. S. Wilkes, and K. E. Johnson, in *Molten Salts XIII*, H. C. Delong, et al., Editors, PV 2002-19, p. 964, The Electrochemical Society Proceedings Series, Pennington, NJ (2002).
7. J. A. Plambeck, *Encyclopedia of Electrochemistry of the Elements*, Vol. X: *Fused Salt Systems*, Marcel Dekker, New York (1976); G. J. Janz, *Molten Salts Handbook*, Academic Press, New York (1967). Both give considerable amounts of data regarding high temperature systems.
8. P. C. Trulove and R. C. Mantz, in *Ionic Liquids in Synthesis*, P. Wasserscheid and T. Welton, Editors, Wiley-VCH, Weinheim, p. 112 (2003). Data are tabulated, with references for more than 500 systems at temperatures of 20-35°C.
9. T. O'Donnell, *Superacids and Acidic Melts*, VCH, Weinheim, p. 8 (1993).
10. I. M. Kolthoff and A. Willman, *J. Am. Chem. Soc.*, **56**, 1007 (1934).

11. P. Vanysek, *CRC Handbook of Chemistry and Physics*, 82<sup>nd</sup> Edition, CRC Press, Boca Raton (2001-2002).
12. J. L. E. Campbell and K. E. Johnson, *J. Am. Chem. Soc.*, **117**, 7791 (1995).
13. G. Driver and K. E. Johnson, *Green Chem.*, **5**, 163 (2003).
14. K. E. Johnson and G. Driver, *Proc. Int. Symp. on Ionic Liquids, in Honor of M. Gaume Escard*, Norwegian Univ. Sci. Tech., Trondheim, p. 233 (2003).
15. G. P. Smith, A. S. Dworkin, R. M. Pagni, and S. P. Zingg, *J. Am. Chem. Soc.*, **111**, 5075 (1989).
16. K. E. Johnson, L. Xiao, and G. Driver, in *ACS Symposium 818*, R. D. Rogers and K. R. Seddon, Editors, Washington, DC, p. 230 (2002).
17. L. Xiao and K. E. Johnson, *J. Electrochem. Soc.*, **150**, E307 (2003).
18. J. A. Gardner, *Ber. Dtsch. Chem. Ges.*, **23**, 1587 (1890).
19. J. von Zawidzki, *Z. Phys. Chem.*, **35**, 129 (1900).
20. I. M. Kolthoff and S. Bruckenstein, *J. Am. Chem. Soc.*, **78**, 1 (1956).
21. G. Charlot and B. Tremillon, *Chemical Reactions in Solvents and Melts*, Pergamon, Oxford (1969).
22. H. A. Laitinen, *Chemical Analysis*, McGraw Hill, New York, p. 57 (1960).
23. R. G. Treble, K. E. Johnson, and E. Tosh, *Can. J. Chem.*, **84**, 915 (2006).
24. M. J. Earle, J. M. S. S. Esperanca, M. A. Gilea, J. N. Canongia Lopes, L. P. N. Rebelo, J. W. Magee, K. R. Seddon, and J. A. Widegren, *Nature*, **439**, 831 (2006).
25. M. Yoshizawa, W. Xu, and C.A. Angell, *J. Am. Chem. Soc.*, **125**, 15411 (2003).

#### About the Author

**KEITH E. JOHNSON** is a Professor Emeritus of Chemistry and a Research Fellow in Energy and Environment at the University of Regina, Canada. A graduate of Imperial College, London, he has worked with ionic liquids of various types for 50 years. He is the author or co-author of over 100 papers, mostly in this field and has directed the research of 37 graduate students and postdoctoral fellows. He joined the ECS in 1960. He may be reached at: Keith.Johnson@uregina.ca.

# THE ELECTROCHEMICAL SOCIETY MONOGRAPH SERIES

The following volumes are sponsored by ECS, and published by John Wiley & Sons, Inc. They should be ordered from: ECS, 65 South Main St., Pennington, NJ 08534-2839, USA.

## Just Published!

**Fundamentals of Electrochemical Deposition (2nd Edition)**  
Edited by M. Paunovic and M. Schlesinger (2006)  
373 pages. ISBN 0-471-71221-3.

**Fundamentals of Electrochemistry (2nd Edition)**  
Edited by V. S. Bagotsky (2005)  
722 pages. ISBN 0-471-70058-4.

**Electrochemical Systems (3rd edition)**  
by John Newman and Karen E. Thomas-Alyea (2004)  
647 pages. ISBN 0-471-47756-7.

**Modern Electroplating (4th edition)**  
by M. Schlesinger and M. Paunovic (2000)  
832 pages. ISBN 0-471-16824-6.

**Atmospheric Corrosion (2nd edition)**  
by C. Leygraf and T. Graedel (2000)  
354 pages. ISBN 0-471-37219-6.

**Uhlig's Corrosion Handbook (2nd edition)**  
by R. Winston Revie (2000)  
1300 pages. ISBN 0-471-15777-5.

**Semiconductor Wafer Bonding**  
by Q. -Y. Tong and U. Gösele (1998)  
297 pages. ISBN 0-471-57481-3.

**Fundamentals of Electrochemical Deposition**  
by M. Paunovic and M. Schlesinger (1998)  
301 pages. ISBN 0-471-16820-3.

**Corrosion of Stainless Steels (2nd edition)**  
by A. J. Sedriks (1996)  
437 pages. ISBN 0-471-00792-7.

**Synthetic Diamond: Emerging CVD Science and Technology**  
Edited by K. E. Spear and J. P. Dismukes (1994)  
688 pages. ISBN 0-471-53589-3.

**Electrochemical Oxygen Technology**  
by K. Kinoshita (1992)  
444 pages. ISBN 0-471-57043-5.

ECS Members will receive a discount. Invoices for the cost of the books plus shipping and handling will be sent after the volumes have been shipped. All prices subject to change without notice.